

Ciprofloxacin -Tablets

Ciprofloxacin -Tablets			
Strength	Composition	Therapeutic Segment	Class
250 mg	Ciprofloxacin HCl BP Eq. to Ciprofloxacin 250 mg	Antibiotic	Fluoroquinolone
500 mg	Ciprofloxacin HCl BP Eq. to Ciprofloxacin 500 mg		
<p>Indications: UTI, Acute Uncomplicated Cystitis in females, Chronic Bacterial Prostatitis, LRTI, Acute Sinusitis, SSTI, Bone and Joint Infections, Complicated Intra-Abdominal Infections, Infectious Diarrhea, Typhoid Fever (Enteric Fever)</p>			
<p>Side-effects: Nausea , diarrhea, liver function tests abnormal, vomiting and rash</p>			
<p>Precautions: Epilepsy, CNS disorder, renal or hepatic dysfunction, G6PD deficiency, good hydration required myasthenia gravis, may impair ability to drive or operate machinery</p>			
<p>Contra-indications: Children < 12 years, Pregnancy, lactation, avoid exposure to sunlight or sun lamps</p>			
<p>Overdose: stomach should be emptied by inducing vomiting or by gastric lavage. supportive treatment, Adequate hydration</p>			
<p>How to take: Do not take with dairy products such as milk . Take this medication for as many days as it has been prescribed</p>			