

Co-Trimoxazole Tablets

Strength	Composition	Therapeutic Segment	Class
480 mg	Trimethoprim BP 80 mg Sulphamethoxazole BP 400 mg	Antibiotic	Dihydrofolate reductase inhibitor/ Sulfonamide antibiotic
960 mg	Trimethoprim BP 160 mg Sulphamethoxazole BP 800 mg		
<p>Indications: Urinary Tract Infections, Acute Otitis Media, Acute Exacerbations of Chronic Bronchitis in Adults, Travelers' Diarrhea in Adults, Shigellosis, Pneumocystis Carinii Pneumonia</p>			
<p>Side-effects: Gastrointestinal disturbances (nausea, vomiting, anorexia) and allergic skin reactions (such as rash and urticaria).</p>			
<p>Precautions: G6PD deficiency, folate deficiency, hepatic and renal impairment, elderly, porphyria, thyroid dysfunction, maintain adequate hydration</p>			
<p>Contra-indications: Hypersensitivity, severe hepatic or renal insufficiency, infants < 2 months, Pregnancy, Lactation</p>			
<p>Overdose: Gastric lavage or emesis, intravenous fluids, Acidification of the urine, blood counts and blood chemistries</p>			
<p>How to take: Take co-trimoxazole with food.</p>			