

## Doxycycline Capsule

Strength	Composition	Therapeutic Segment	Class
100 mg	Doxycycline Hydrochloride BP 100 mg	Antibiotic	Tetracycline
<p><b>Indications:</b> Rocky mountain spotted fever, typhus fever and the typhus group, Q fever, rickettsialpox, and tick fevers Respiratory tract infections, Lymphogranuloma venereum, Psittacosis, Trachoma, Inclusion conjunctivitis etc</p>			
<p><b>Side-effects:</b> Anorexia, nausea, vomiting, diarrhea, glossitis, dysphagia, enterocolitis, and inflammatory lesions Photosensitivity, Hypersensitivity Reactions, Bulging fontanels in infants and intracranial hypertension in adults</p>			
<p><b>Precautions:</b> Impaired hepatic function, oral candidiasis, should be taken in upright position to avoid esophageal injury</p>			
<p><b>Contra-indications:</b> child &lt; 8yrs, pregnancy, lactation, hypersensitivity to tetracycline, prolonged exposure to sunlight</p>			
<p><b>Overdose:</b> discontinue medication, treat symptomatically and institute supportive measures.</p>			
<p><b>How to take:</b> Drink a full glass of water with each dose. Continue to take doxycycline even if you feel well</p>			